


Directions for Use

Step-by-step instructions for taking RELYVRIO™

 **relyvrio™**
(sodium phenylbutyrate and
taurursodiol) for oral
suspension 3 g/1 g

This quick overview offers general recommendations about how to take RELYVRIO. Be sure to speak with your doctor, who is always the best source for treatment information, before beginning RELYVRIO.



Recommended dosing for RELYVRIO

For the first 3 weeks: Take just 1 packet of RELYVRIO every day.

After 3 weeks: Take 2 packets of RELYVRIO per day, taking 1 packet in the morning and 1 packet at night.

Starting with 1 packet a day for the first 3 weeks may help improve tolerability.

Additional dosing information

Avoid taking antacids that contain aluminum. RELYVRIO should be taken before a snack or meal.

What to do if you miss a dose

If you forget to take one or more doses, you should take your next dose at the normal time and in the normal amount. You should not take more than what is prescribed by your doctor.

INDICATION


RELYVRIO is indicated for the treatment of amyotrophic lateral sclerosis (ALS) in adults.

For more information about RELYVRIO, please see the Important Safety Information on the back and the full [Prescribing Information and Medication Guide](#).


HOW TO TAKE RELYVRIO



Use scissors to open the packet of RELYVRIO.



Pour contents of packet into a cup.



Add approximately 1 cup (8 oz) of room temperature water.

Do not mix RELYVRIO with liquid other than water.^a



Stir vigorously until dissolved.

This may require significant stirring or gentle crushing.



Take each dose of RELYVRIO in its entirety within 1 hour of dissolving.

You can drink it or take it via a feeding tube.




**Optional for feeding tube administration:*

Use a clean gravity bag or syringe to administer RELYVRIO through the feeding tube.

After each dose, it's important to flush the feeding tube with warm water, as directed by your doctor.

^aWhen mixed, the solution may appear cloudy and/or contain residue.

Additional information you should know

 **relyvrio™**
(sodium phenylbutyrate and
taurursodiol) for oral
suspension 3 g/1 g



RELYVRIO™ AND OTHER TREATMENTS

RELYVRIO may be taken alone or with other FDA-approved treatments for ALS. Before taking any additional medications, talk to your doctor.



STORAGE

RELYVRIO should be stored at room temperature (between 68° F and 77° F) and protected from moisture. Keep out of reach and sight of children.



A SUPPORT PROGRAM YOU CAN COUNT ON

The Amylx Care Team (ACT)™ is a personalized support program designed to help meet your individual needs as you start and continue on RELYVRIO. Along with your doctor and specialty pharmacy, ACT team members can help you with any questions you might have about taking RELYVRIO.

Learn more and get in touch with ACT today by calling **866-318-2989**, visiting [AmylxCareTeam.com](https://www.amylix.com/care-team), or scanning this QR code.



IMPORTANT SAFETY INFORMATION

Before you receive RELYVRIO, tell your doctor about all of your medical conditions, including if you:

- Have pancreas, liver, or intestinal problems
- Have heart failure, including congestive heart failure
- Have high blood pressure
- Have kidney problems
- Are pregnant or plan to become pregnant. It is not known if RELYVRIO will harm your unborn baby
- Are breastfeeding or plan to breastfeed. It is not known if RELYVRIO passes into your breast milk. You and your doctor should decide the best way to feed your baby

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements and any taurursodiol products, such as taurursodeoxycholic acid (TUDCA).

RELYVRIO may affect the way other medicines work, and other medicines may affect how RELYVRIO works.

What are the possible side effects of RELYVRIO?

RELYVRIO may cause serious side effects, including:

- **Changes in bile acid levels.** RELYVRIO may increase bile acid levels and cause

worsening diarrhea if you already have problems with your liver, bile ducts, or pancreas. Your doctor should monitor you for these side effects. Some disorders of the pancreas, bile ducts, or intestines may also make it harder to absorb RELYVRIO

- **Salt (sodium) retention.** RELYVRIO contains a high amount of salt. For people who are sensitive to salt intake, such as people with heart failure, high blood pressure, or kidney problems, limit the amount of salt you eat and drink. Talk to your doctor about the total amount of daily salt that is right for you. Your doctor will monitor you for signs and symptoms of salt retention during your treatment with RELYVRIO

The most common side effects of RELYVRIO include diarrhea, abdominal pain, nausea, and upper respiratory tract infection.

Tell your doctor if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of RELYVRIO. Call your doctor for medical advice about side effects.

You may report side effects to FDA at [1-800-FDA-1088](https://www.fda.gov/medwatch).

For more information about RELYVRIO, please see the full Prescribing Information and Medication Guide.



©2022 Amylx Pharmaceuticals, Inc. All Rights Reserved. AMYLYX®, the AMYLYX® logo, RELYVRIO™, the RELYVRIO™ logo, and AMYLYX CARE TEAM (ACT)™ are trademarks of Amylx Pharmaceuticals, Inc. PM-00143 09/22