

“LET’S GET STARTED”

Tyrone
Living with ALS

Useful information to help you start with confidence on your RELYVRIO™ (sodium phenylbutyrate and taurursodiol) treatment journey

WHAT TO EXPECT

RELYVRIO is a new prescription oral therapy for adults living with amyotrophic lateral sclerosis (ALS) shown to help slow disease progression.



RELYVRIO IS NOT A CURE

While RELYVRIO will not stop ALS, it can help slow down disease progression.



WHAT TO KNOW ABOUT SIDE EFFECTS

In the CENTAUR clinical trial for RELYVRIO, the most common side effects were diarrhea, abdominal pain, nausea, and upper respiratory tract infection. Gastrointestinal side effects occurred throughout the study but were more frequent during the first 3 weeks of treatment.



RELYVRIO AND OTHER TREATMENTS

RELYVRIO may be taken alone or with other FDA-approved treatments for ALS. Before taking any additional medications, talk to your doctor.

IMPORTANT SAFETY INFORMATION

Before you receive RELYVRIO, tell your doctor about all of your medical conditions, including if you:

- Have pancreas, liver, or intestinal problems
- Have heart failure, including congestive heart failure
- Have high blood pressure
- Have kidney problems

(continued on back)


For more information about RELYVRIO, please see the additional Important Safety Information on the back and the full [Prescribing Information and Medication Guide](#).



MANAGING THE TASTE

The taste of RELYVRIO can vary from person to person.

- When mixed with water, RELYVRIO may have a fruity smell, but that may not reflect its actual taste
- RELYVRIO may have a bitter aftertaste. In the clinical trial, participants were allowed to try to lessen or remove the bitter aftertaste by:
 - using mint-flavored mouth strips or mouth spray immediately before or after taking the medicine
 - eating a small snack or honey after taking RELYVRIO
 - drinking milk after taking RELYVRIO
 - avoiding fruit juice immediately afterwards, as it can make the aftertaste worse

 **relyvrio™**
(sodium phenylbutyrate and taurursodiol) for oral suspension 3 g/1 g

THE AMYLYX CARE TEAM (ACT)[™] IS HERE TO HELP



Our team can help with the process of starting and continuing on RELYVRIO[™] (sodium phenylbutyrate and taurursodiol)

IS RELYVRIO RIGHT FOR YOU?

First, talk to your doctor to determine if RELYVRIO is right for you. It's important to reach out to them and start the enrollment process as soon as possible. **It may take some time between submitting the RELYVRIO Enrollment Form and getting RELYVRIO delivered to your home.** ACT team members can explain the process and keep you updated on important milestones.



Learn more and get in touch with ACT today by calling **866-318-2989**, visiting **[AmylyxCareTeam.com](https://www.AmylyxCareTeam.com)**, or scanning this **QR code**.

IMPORTANT SAFETY INFORMATION

(continued from front)

- Are pregnant or plan to become pregnant. It is not known if RELYVRIO will harm your unborn baby
- Are breastfeeding or plan to breastfeed. It is not known if RELYVRIO passes into your breast milk. You and your doctor should decide the best way to feed your baby

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements and any taurursodiol products, such as taurursodeoxycholic acid (TUDCA).

RELYVRIO may affect the way other medicines work, and other medicines may affect how RELYVRIO works.

What are the possible side effects of RELYVRIO?

RELYVRIO may cause serious side effects, including:

- **Changes in bile acid levels.** RELYVRIO may increase bile acid levels and cause worsening diarrhea if you already have problems with your liver, bile ducts, or pancreas. Your doctor should monitor you for these side effects. Some disorders of the pancreas, bile ducts, or intestines may also make it harder to absorb RELYVRIO
- **Salt (sodium) retention.** RELYVRIO contains a high amount of salt. For people who are sensitive to salt intake, such as people with heart failure, high blood pressure, or kidney problems, limit the amount of salt you eat and drink. Talk to your doctor about the total amount of daily salt that is right for you. Your doctor will

monitor you for signs and symptoms of salt retention during your treatment with RELYVRIO

The most common side effects of RELYVRIO include diarrhea, abdominal pain, nausea, and upper respiratory tract infection.

Tell your doctor if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of RELYVRIO. Call your doctor for medical advice about side effects.

You may report side effects to FDA at [1-800-FDA-1088](https://www.fda.gov/medwatch).

INDICATION

RELYVRIO is indicated for the treatment of amyotrophic lateral sclerosis (ALS) in adults.

For more information about RELYVRIO, please see the full [Prescribing Information and Medication Guide](#).



©2022 Amylyx Pharmaceuticals, Inc. All Rights Reserved. AMYLYX[®], the AMYLYX[®] logo, RELYVRIO[™], the RELYVRIO[™] logo, AMYLYX CARE TEAM (ACT)[™], and the AMYLYX CARE TEAM (ACT)[™] logo are trademarks of Amylyx Pharmaceuticals, Inc. PM-00059 09/22

